

PRODUCT	CODE	QTY	USD
Resins			
<i>o</i>-BAL-CLEAR Resin <i>ortho</i> -Backbone Amide Linker attached to CLEAR Resin for the synthesis of C-terminally Modified Peptides (peptide acids, esters, aldehydes, alcohols, head-to-tail cyclic peptides)	CBL-1291-PI 4 °C	Please Inquire	
<i>p</i>-BAL-CLEAR Resin <i>para</i> -Backbone Amide Linker attached to CLEAR Resin for the synthesis of C-terminally Modified Peptides (peptide acids, esters, aldehydes, alcohols, head-to-tail cyclic peptides)	CBL-1292-PI 4 °C	Please Inquire	
Fmoc-D-Lys(Boc)-CLEAR-Acid Resin (100-200 mesh) 1% DVB	CFK-1273-PI 4 °C	1 g 5 g	120 435
Fmoc-D-Ala-Wang Resin (100-200 mesh) 1% DVB	RFA-1356-PI 4 °C	1 g 5 g	60 240
Fmoc-D-Arg(Pbf)-Wang Resin (100-200 mesh) 1% DVB	RFR-1347-PI 4 °C	1 g 5 g	60 240
Fmoc-D-Leu-Wang Resin (100-200 mesh) 1% DVB	RFL-1357-PI 4 °C	1 g 5 g	60 240
H-Asn-2-Citrt-Resin (100-200 mesh) 1% DVB	RHN-11052-PI 4 °C	1 g 5 g	30 115
H-Gln-2-Citrt-Resin (100-200 mesh) 1% DVB	RHQ-11057-PI 4 °C	1 g 5 g	30 115
H-mini-PEG-2-CITrt-Resin (100-200 mesh) 1% DVB mini-PEG™ = 8-Amino-3,6-Dioxaoctanoic Acid	RHX-11074-PI 4 °C	1 g 5 g	99 396
TentaGel S PHB-Gly-Fmoc Resin (90 μm)	RTS-9950-PI 4 °C	1 g 5 g	59 214
TentaGel S PHB-Leu-Fmoc Resin (90 μm)	RTS-9951-PI 4 °C	1 g 5 g	59 214
Boc-Lys-(Fmoc)-O-Resin (100-200 mesh) 1% DVB	RBK-1139-PI 4 °C	1 g 5 g	75 275
Reagents			
<i>o</i>-BAL 5-(2-Formyl-3,5-Dimethoxyphenoxy)Pentanoic Acid <i>ortho</i> -Backbone Amide Linker (M.W. 282.30) C ₁₄ H ₁₆ O ₆ Handle for the Synthesis of C-terminally Modified Peptides (peptide acids, esters, aldehydes, alcohols, head-to-tail cyclic peptides)	KBL-1002-PI -20 °C	5 g 25 g	235 940
<i>p</i>-BAL 5-(4-Formyl-3,5-Dimethoxyphenoxy)Pentanoic Acid <i>para</i> -Backbone Amide Linker (M.W. 282.30) C ₁₄ H ₁₈ O ₆ Handle for the Synthesis of C-terminally Modified Peptides (peptide acids, esters, aldehydes, alcohols, head-to-tail cyclic peptides)	KBL-1006-PI -20 °C	5 g 25 g	235 940
Biologically Active Peptides			
Arg-Gly-Asp (RGD) Related Peptides			
DOTA-Glu-[<i>cyclo</i> (Arg-Gly-Asp-D-Phe-Lys)]₂ DOTA-E-[α(RGDfK)]₂ (M.W. 1704.88) C ₇₅ H ₁₁₃ N ₂₃ O ₂₃ α ₃ Integrin Binding RGD Peptide RGD Tumor Targeting Peptide for Radiolabelling (Requires further derivatization before use)	PCI-3903-PI -20 °C	1 mg 5 mg	89 345
M.L. Janssen, W.J. Oyen, I. Dijkgraaf, L.F. Massuger, C. Frielink, D.S. Edwards, M. Rajopadhye, H. Boonstra, F.H. Corstens, and O.C. Boerman, <i>Cancer Res.</i> , 62 , 6146 (2002). X. Chen, S. Liu, Y. Hou, M. Tohme, R. Park, J.R. Bading, and P.S. Conti, <i>Mol. Imaging Biol.</i> , 6 , 350 (2004).			
H-Glu[<i>cyclo</i> (Arg-Gly-Asp-D-Phe-Lys)]₂ H-E[α(RGDfK)]₂ Trifluoroacetate Salt (M.W. 1318.47) C ₅₉ H ₈₇ N ₁₉ O ₁₆ RGD Tumor Targeting and Tumor Imaging Peptide (Requires further derivatization before use).	PCI-3904-PI -20 °C	1 mg 5 mg 25 mg	69 225 795
H-Gly-Arg-Gly-Glu-Ser-OH Fibronectin Active Fragment Negative Control	PFA-3907-PI -20 °C	5 mg 25 mg	50 225

PRODUCT	CODE	QTY	USD
H-Gly-Arg-Ala-Asp-Ser-Pro-OH GRADSP	PCI-3910-PI -20 °C	1 mg 5 mg	45 175
H-Gly-Arg-Gly-Asp-Asn-Pro-OH GRGDNP	PCI-3909-PI -20 °C	5 mg 25 mg	75 295
H-Glu[<i>cyclo</i> (Arg-Gly-Asp-D-Tyr-Lys)]₂ H-E[α (RGDyK)] ₂ (M.W. 1350.47) C ₅₉ H ₈₇ N ₁₉ O ₁₈ <i>RGD Tumor Targeting and Tumor Imaging Peptide</i> (Requires further derivatization before use) Y. Wu, X. Zhang, Z. Xiong, Z. Cheng, D.R. Fisher, S. Liu, S.S. Gambhir, and X. Chen, <i>J. Nucl. Med.</i> , 46 , 1707 (2005). X. Chen, S. Liu, Y. Hou, M. Tohme, R. Park, J.R. Bading, and P.S. Conti, <i>Mol. Imaging Biol.</i> , 6 , 350 (2004). X. Chen, C. Plasencia, Y. Hou, and N. Neamati, <i>J. Med. Chem.</i> , 48 , 1098 (2005). X. Chen, M. Tohme, R. Park, Y. Hou, J.R. Bading, and P.S. Conti, <i>Mol. Imaging</i> , 3 , 96 (2004).	PCI-3899-PI -20 °C	1 mg 5 mg 25 mg	69 225 795
<i>cyclo</i> (Arg-Ala-Asp-D-Tyr-Lys) α (RADyK) (M.W. 633.71) C ₂₈ H ₄₃ N ₉ O ₈ <i>Control Peptide of cyclo (Arg-Gly-Asp-D-Tyr-Lys) (PCI-3662-PI)</i>	PCI-3894-PI -20 °C	1 mg 5 mg 25 mg	49 149 595
<i>cyclo</i> [Arg-Gly-Asp-D-Phe-Lys(Biotin)] α (RGDfK(Biotin)) Acetate Salt (M.W. 829.98) C ₃₇ H ₅₅ N ₁₁ O ₉ S α , β ₃ <i>Integrin Binding RGD Peptide RGD Tumor Targeting Peptide</i> (Requires further derivatization before use)	PCI-3895-PI -20 °C	1 mg 5 mg	59 175
<i>cyclo</i> [Arg-Gly-Asp-D-Phe-Lys(H-Ser)] α (RGDfK(H-Ser)) (M.W. 690.76) C ₃₀ H ₄₆ N ₁₀ O ₉ α , β ₃ <i>Integrin Binding RGD Peptide RGD Tumor Targeting Peptide</i> (Requires further derivatization before use)	PCI-3898-PI -20 °C	1 mg 5 mg	59 175
Defensin Peptides			
α-Defensin-2 (Human) <i>Antimicrobial Peptide</i>	PDF-4428-s -20 °C	0.1 mg vial	212
Ghrelin Related Peptides			
[Trp³, Arg⁵]-Ghrelin (1-5) GSWFR H-Gly-Ser-Trp-Phe-Arg-OH (M.W. 651.73) C ₃₁ H ₄₁ N ₉ O ₇ <i>Growth-Hormone Secretagogue (GHS) Receptor Agonist / Stimulates Food-Intake</i> K. Ohinata, K. Kobayashi, M. Yoshikawa, <i>Peptides</i> , 27 , 1632 (2006).	PGH-3902-PI -20 °C	1 mg 5 mg	35 95
Des-Acyl Ghrelin (Human) Des-<i>n</i>-Octanoyl Ghrelin (Human) Trifluoroacetate Salt Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Arg-Val-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (M.W. 3244.7) C ₁₄₁ H ₂₃₅ N ₄₇ O ₄₁ <i>Des-Octanoylated Ghrelin with Distinct Effect on Food Intake</i> • This compound is distributed through Peptide Institute, Inc. under the license of Dr. K. Kangawa.	PGH-4436-s -20 °C	0.1 mg vial	115
Des-Acyl Ghrelin (Rat) Des-<i>n</i>-Octanoyl Ghrelin (Rat) Trifluoroacetate salt Gly-Ser-Ser-Phe-Leu-Ser-Pro-Glu-His-Gln-Lys-Ala-Gln-Gln-Arg-Lys-Glu-Ser-Lys-Lys-Pro-Pro-Ala-Lys-Leu-Gln-Pro-Arg (M.W. 3188.6) C ₁₃₉ H ₂₃₁ N ₄₅ O ₄₁ <i>Des-Octanoylated Ghrelin with Distinct Effect on Food Intake</i> • This compound is distributed through Peptide Institute, Inc. under the license of Dr. K. Kangawa.	PGH-4437-s -20 °C	0.1 mg vial	115
Guangxitoxin			
Guangxitoxin-1E <i>Kv2.1/Kv2.2 Channel Blocker / Enhancer of Glucose-Stimulated Insulin Secretion</i> J. Herrington, Y.-P. Zhou, R.M. Bugianesi, P.M. Dulski, Y. Feng, V.A. Warren, M.M. Smith, M.G. Kohler, V.M. Garsky, M. Sanchez, M. Wagner, K. Raffaelli, P. Banerjee, C. Ahaghotu, D. Wunderler, B.T. Priest, J.T. Mehl, M.L. Garcia, O.B. McManus, G.J. Kaczorowski, and R.S. Slaughter, <i>Diabetes</i> , 55 , 1034-1042 (2006).	PGX-4433-s -20 °C	0.1 mg vial	229

PRODUCT	CODE	QTY	USD
Liver-Expressed Antimicrobial Peptides			
Hepcidin (Mouse) <i>(Coming Soon)</i> Liver-Expressed Antimicrobial Peptide 1 (LEAP-1) (Mouse) <i>Iron-Regulatory Hormone</i>	PLP-4434-s -20 °C	0.1 mg	229
Melanotropins			
H-Tyr-Val-Met-Gly-His-Phe-Arg-D-Trp-Asp-Arg-Phe-Gly-OH YVMGHFRwDRFG (M.W. 1570.81) C ₇₄ H ₉₉ N ₂₁ O ₁₆ S <i>Potent and Selective Melanocortin 3 (MC3) Receptor Agonist</i>	PMC-3897-PI -20 °C	1 mg 5 mg	49 195
D. Marks, V.J. Hruby, G. Brookhart, and R.D. Cone, <i>Peptides</i> , 27 , 259 (2006). P. Grieco, P.M. Balse, D. Weinberg, T. MacNeil, and V.J. Hruby, <i>J. Med. Chem.</i> , 43 , 4998 (2000).			
Muscarinic Toxins			
Muscarinic Toxin 3 [MT3, MTX3, m4-toxin] (Green Mamba, <i>Dendroaspis angusticeps</i>) Synthetic Product Leu-Thr-Cys-Val-Thr-Lys-Asn-Thr-Ile-Phe-Gly-Ile-Thr-Thr-Glu-Asn-Cys-Pro-Ala-Gly-Gln-Asn-Leu-Cys-Phe-Lys-Arg-Trp-His-Tyr-Val-Ile-Pro-Arg-Tyr-Thr-Glu-Ile-Thr-Arg-Gly-Cys-Ala-Ala-Thr-Cys-Pro-Ile-Pro-Glu-Asn-Tyr-Asp-Ser-Ile-His-Cys-Cys-Lys-Thr-Asp-Lys-Cys-Asn-Glu (Disulfide bonds between Cys ³ -Cys ²⁴ , Cys ¹⁷ -Cys ⁴² , Cys ⁴⁶ -Cys ⁵⁷ , and Cys ⁵⁸ -Cys ⁶³) (M.W. 7379.4) C ₃₁₉ H ₄₈₉ N ₈₉ O ₉₇ S ₈ <i>Specific Ligand for Muscarinic Acetylcholine Receptor-4 (M₄)</i>	PMT-4410-s -20 °C	0.1 mg vial	315
M. Jolkkonen, P.L.M. van Giersbergen, U. Hellman, C. Wernstedt, and E. Karlsson, <i>FEBS Lett.</i> , 352 , 91 (1994). (Original; MT3) J.-S. Liang, J. Carsi-Gabrenas, J.L. Krajewski, J.M. McCafferty, S.L. Purkerson, M.P. Santiago, W.L. Strauss, H.H. Valentine, and L.T. Potter, <i>Toxicon</i> , 34 , 1257 (1996). (Original; m4-toxin) A. Adem and E. Karlsson, <i>Life Sci.</i> , 60 , 1069 (1997). (<i>Pharmacol.; Muscarinic Receptor Subtype Specificity</i>) S. Katayama, M. Ishimaru, H. Nishio, Y. Nishiuchi, and T. Kimura, <i>Peptide Science</i> 2004, 161 (2005). (S-S Bond)			
Muscarinic Toxin α <i>(Coming Soon)</i> MTα (Black Mamba, <i>Dendroaspis polylepis</i>) Leu-Thr-Cys-Val-Thr-Ser-Lys-Ser-Ile-Phe-Gly-Ile-Thr-Thr-Glu-Asn-Cys-Pro-Asp-Gly-Gln-Asn-Leu-Cys-Phe-Lys-Lys-Trp-Tyr-Tyr-Leu-Asn-His-Arg-Tyr-Ser-Asp-Ile-Thr-Trp-Gly-Cys-Ala-Ala-Thr-Cys-Pro-Lys-Pro-Thr-Asn-Val-Arg-Glu-Thr-Ile-His-Cys-Cys-Glu-Thr-Asp-Lys-Cys-Asn-Glu (Disulfide bonds between Cys ³ -Cys ²⁴ , Cys ¹⁷ -Cys ⁴² , Cys ⁴⁶ -Cys ⁵⁸ , and Cys ⁵⁹ -Cys ⁶⁴) (M.W. 7545.4) C ₃₂₆ H ₄₉₁ N ₈₉ O ₁₀₂ S ₈ <i>Ligand for Muscarinic Acetylcholine Receptor-3/5 (M3/M5) (Non-specific Ligand)</i>	PMT-4424-s -20 °C	0.1 mg vial	315
M. Jolkkonen, P.L.M. van Giersbergen, U. Hellman, C. Wernstedt, A. Oras, N. Satyapan, A. Adem, and E. Karlsson, <i>Eur. J. Biochem.</i> , 234 , 579 (1995). (Original; MTα) A. Adem and E. Karlsson, <i>Life Sci.</i> , 60 , 1069 (1997). (<i>Pharmacol.; Muscarinic Receptor Subtype Specificity</i>) M. Jolkkonen, A. Oras, T. Toomela, E. Karlsson, J. Järvi, and K.E.O. Åkerman, <i>Toxicon</i> , 39 , 377 (2001). (<i>Pharmacol.; Mechanism of Receptor Binding</i>)			
Neuromedins			
Neuromedin U (Porcine, 1-8) H-Tyr-Phe-Leu-Phe-Arg-Pro-Arg-Asn-NH ₂ (M.W. 1111.32) C ₅₄ H ₇₈ N ₁₆ O ₁₀ <i>Neuropeptide</i>	PNM-3698-PI -20 °C	1 mg 5 mg	35 139
Neuromedin S (Human) NMS (Human) Ile-Leu-Gln-Arg-Gly-Ser-Gly-Thr-Ala-Ala-Val-Asp-Phe-Thr-Lys-Lys-Asp-His-Thr-Ala-Thr-Trp-Gly-Arg-Pro-Phe-Phe-Leu-Phe-Arg-Pro-Arg-Asn-NH ₂ (M.W. 3791.3) C ₁₇₃ H ₂₆₅ N ₅₃ O ₄₄ <i>Food Intake Suppressor / Regulator of Circadian Rhythm</i>	PNM-4426-s -20 °C	1 mg vial	116
K. Mori, M. Miyazato, T. Ida, N. Murakami, R. Serino, Y. Ueta, M. Kojima, and K. Kangawa, <i>EMBO J.</i> , 24 , 325 (2005). (Original) T. Ida, K. Mori, M. Miyazato, Y. Egi, S. Abe, K. Nakahara, M. Nishihara, K. Kangawa, and N. Murakami, <i>Endocrinology</i> , 146 , 4217 (2005). (<i>Pharmacol.; Anorexigenic Hormone</i>)			
• This compound is distributed through Peptide Institute, Inc. under the license of National Cardiovascular Center and Takeda Chemical Industries, Ltd.			

PRODUCT	CODE	QTY	USD
Neuromedin S (Rat) NMS (Rat) Leu-Pro-Arg-Leu-Leu-His-Thr-Asp-Ser-Arg-Met-Ala-Thr-Ile-Asp-Phe-Pro-Lys-Lys-Asp-Pro-Thr-Thr-Ser-Leu-Gly-Arg-Pro-Phe-Phe-Leu-Phe-Arg-Pro-Arg-Asn-NH ₂ (M.W. 4241.9) C ₁₉₃ H ₃₀₇ N ₅₇ O ₄₉ S <i>Food Intake Suppressor / Regulator of Circadian Rhythm</i> K. Mori, M. Miyazato, T. Ida, N. Murakami, R. Serino, Y. Ueta, M. Kojima, and K. Kangawa, <i>EMBO J.</i> , 24 , 325 (2005). (Original) T. Ida, K. Mori, M. Miyazato, Y. Egi, S. Abe, K. Nakahara, M. Nishihara, K. Kangawa, and N. Murakami, <i>Endocrinology</i> , 146 , 4217 (2005). (Pharmacol.; Anorexigenic Hormone) • This compound is distributed through Peptide Institute, Inc. under the license of National Cardiovascular Center and Takeda Chemical Industries, Ltd.	PNM-4427-s -20 °C	0.1 mg vial	126
Obestatin			
Obestatin (Human) Phe-Asn-Ala-Pro-Phe-Asp-Val-Gly-Ile-Lys-Leu-Ser-Gly-Val-Gln-Tyr-Gln-Gln-His-Ser-Gln-Ala-Leu-NH ₂ (M.W. 2546.8) C ₁₁₆ H ₁₇₆ N ₃₂ O ₃₃ <i>Food Intake Suppressor / Ligand for GPR39</i> J.V. Zhang, P.-G. Ren, O. Avsian-Kretchmer, C.-W. Luo, R. Rauch, C. Klein, and A.J.W. Hsueh, <i>Science</i> , 310 , 996 (2005). (Original; Structure & Pharmacol.)	PGH-4429-s -20 °C	0.1 mg vial	69
Obestatin (Rat, Mouse) Phe-Asn-Ala-Pro-Phe-Asp-Val-Gly-Ile-Lys-Leu-Ser-Gly-Ala-Gln-Tyr-Gln-Gln-His-Gly-Arg-Ala-Leu-NH ₂ (M.W. 2516.8) C ₁₁₄ H ₁₇₄ N ₃₄ O ₃₁ <i>Food Intake Suppressor / Ligand for GPR39</i> J.V. Zhang, P.-G. Ren, O. Avsian-Kretchmer, C.-W. Luo, R. Rauch, C. Klein, and A.J.W. Hsueh, <i>Science</i> , 310 , 996 (2005). (Original; Structure & Pharmacol.)	PGH-4430-s -20 °C	0.1 mg vial	69
PAR Peptides			
H-Val-Lys-Gly-Ile-Leu-Ser-NH₂ VKGILS-Amide M.W. 614.79 C ₂₈ H ₅₄ N ₈ O ₇ <i>Protease-Activated Receptor 2 (PAR₂) Negative Control Peptide of PAR-3889-PI</i> S.K. Bohm, W. Kong, D. Bromme, S.P. Smekens, D.C. Anderson, A. Connolly, M. Kahn, H.A. Nelken, S.R. Coughlin, D.G. Payan, and N.W. Bunnett, <i>Biochem. J.</i> , 314 , 1009 (1996).	PAR-3896-PI -20 °C	1 mg 5 mg	35 139
H-Tyr-Arg-Leu-Leu-Ser-Phe-NH₂ YRLLSF-Amide (M.W. 796.46) C ₃₉ H ₆₀ N ₁₀ O ₈ <i>Protease-Activated Receptor 2 (PAR₂) Negative Control Peptide of PAR-3888-PI</i>	PAR-3906-PI -20 °C	1 mg 5 mg	35 139
Plectasin			
Plectasin (Coming Soon) <i>Antimicrobial Peptide</i>	PDF-4432-s -20 °C	0.1 mg vial	239
ProTx-I			
ProTx-1 (Tarantula, <i>Thrixopelma pruriens</i>) Glu-Cys-Arg-Tyr-Trp-Leu-Gly-Gly-Cys-Ser-Ala-Gly-Gln-Thr-Cys-Cys-Lys-His-Leu-Val-Cys-Ser-Arg-Arg-His-Gly-Trp-Cys-Val-Trp-Asp-Gly-Thr-Phe-Ser (Disulfide bonds between Cys ² -Cys ¹⁶ , Cys ⁹ -Cys ²¹ , and Cys ¹⁵ -Cys ²⁸) (M.W. 3987.5) C ₁₇₁ H ₂₄₅ N ₅₃ O ₄₇ S ₆ <i>T-Type Ca²⁺ Channel / Na⁺ Channel / K⁺ Channel Blocker (Gating Modifier)</i> R.E. Middleton, V.A. Warren, R.L. Kraus, J.C. Hwang, C.J. Liu, G. Dai, R.M. Brochu, M.G. Kohler, Y.-D. Gao, V.M. Garsky, M.J. Bogusky, J.T. Mehl, C.J. Cohen, and M.M. Smith, <i>Biochemistry</i> , 41 , 14734 (2002). (Original)	PTX-4409-s -20 °C	0.1 mg vial	232
Psalmotoxin 1			
Psalmotoxin 1 (Coming Soon) <i>Selective Blocker for Acid-Sensitive Ion Channel, ASIC1a</i>	PTX-4435-s -20 °C	0.1 mg vial	245
RGD Peptides – See Arg-Gly-Asp Related Peptides			
Substrates			
Abz-Ala-Gly-Leu-Ala-p-Nitro-Benzyl-Amide Trifluoroacetate Salt 2-Aminobenzoyl-L-Alanyl-Glycyl-L-Leucyl-L-Alanyl- <i>para</i> -Nitro-Benzyl-Amide (M.W. 583.65) C ₂₈ H ₃₇ N ₇ O ₇ <i>Substrate for Thermolysin and Neutral Endopeptidase 24.11 (NEP)</i> N. Nishino and J.C. Powers, <i>J. Biol. Chem.</i> , 255 , 3482 (1980). R.S. Rush, M. Mitas, J.C. Powers, T. Tanaka, and L.B. Hersh, <i>Arch. Biochem. Biophys.</i> , 231 , 390 (1984). D.I. Mundy and W.J. Strittmatter, <i>Cell</i> , 40 , 645 (1985)	SAG-3905-PI -20 °C	5 mg 25 mg	45 175

PRODUCT	CODE	QTY	USD
Inhibitors			
Z-Val-Ala-Asp(OMe)-CH₂F [Z-VAD-FMK] {Benzyloxycarbonyl-L-Valyl-L-Alanyl-[(2S)-2-Amino-3-(Methoxycarbonyl)propionyl]} Fluoromethane (M.W. 467.49) C ₂₂ H ₃₀ N ₃ O ₇ F Synthetic Product <i>Inhibitor for Caspases</i>	ICA-3188-v -20 °C	1 mg vial	109
E.A. Slee, H. Zhu, S.C. Chow, M. MacFarlane, D.W. Nicholson, and G.M. Cohen, <i>Biochem. J.</i> , 315 , 21 (1996). H. Zhu, H.O. Fearnhead, and G.M. Cohen, <i>FEBS Lett.</i> , 374 , 303 (1995)			
Pepstatin A (Purity Higher than 90% by HPLC) Microbial Product	IPA-4397 -20 °C	25 mg 100 mg	75 178
In 2003, we began distributing Pepstatin A of “purity higher than 90%” when analyzed by RP-HPLC (code IPA-4397). This new item was welcomed by a majority of our customers. However, this accompanied some complaints because of its low solubility. At the beginning, we thought this low solubility was the destiny of the higher purity Pepstatin A. In striving to offer our customers satisfactory service, we continued to improve the solubility of this item. Recently, we have succeeded in obtaining both higher purity and higher solubility Pepstatin A. This new improved Pepstatin A is now available from us with easier handling.			
Carbohydrates & Conjugates			
Antiproliferative Factor Sialoglycopeptide (see page 221) <i>Antiproliferative Factor from Interstitial Cystitis Patients</i>	CAR-24007-v -20 °C	50 µg vial	212
Peptide Antisera			
Adrenomedullin 2 (Human, 1-7) Antiserum (Rabbit)	NAD-14438-v -20 °C	50 µL vial	339
Antiserum: lyophilized from 0.001M Phosphate Buffer (pH 7.0) Immunogen: Adrenomedullin 2 (Human, 1-7)-TG (TG: Bovine Thyroglobulin)			
Reactivity: Adrenomedullin 2 (Human) + Adrenomedullin (Human) - Calcitonin (Human) - CGRP (Human) -			
Salusin-β (Human) Antiserum (Rabbit)	NLS-14418-v -20 °C	50 µL vial	339
Antiserum: lyophilized from 0.001M Phosphate Buffer (pH 7.0) Immunogen: Salusin β (Human7)-TG (TG: Bovine Thyroglobulin)			
Reactivity: Salusin β (Human) + Salusin α (Human) -			